

Pocket SCAT2



FIFA®

Concussion should be suspected in the presence of **any one or more** of the following: symptoms (such as headache), or physical signs (such as unsteadiness), or impaired brain function (e.g. confusion) or abnormal behaviour.

1. Symptoms

Presence of any of the following signs & symptoms may suggest a concussion.

Loss of consciousness

Feeling slowed down

Seizure or convulsion

Feeling like “in a fog”

Amnesia

“Don’t feel right”

Headache

Difficulty concentrating

“Pressure in head”

Difficulty remembering

Neck Pain

Fatigue or low energy

Nausea or vomiting

Confusion

Dizziness

Drowsiness

Blurred vision

More emotional

Balance problem

Irritability

Sensitivity to light

Sadness

Sensitivity to noise

Nervous or anxious



2. Memory function

Failure to answer all questions correctly may suggest a concussion.

“At what venue are we at today?”

“Which half is it now?”

“Who scored last in this game?”

“What team did you play last week/game?”

“Did your team win the last game?”

3. Balance testing

Instructions for tandem stance

*“Now stand heel-to-toe with your **non-dominant** foot in back. Your weight should be evenly distributed across both feet. You should try to maintain stability for 20 seconds with your hands on your hips and your eyes closed. I will be counting the number of times you move out of this position. If you stumble out of this position, open your eyes and return to the start position and continue balancing. I will start timing when you are set and have closed your eyes.”*

Observe the athlete for 20 seconds. If they make more than 5 errors (such as lift their hands off their hips; open their eyes; lift their fore-foot or heel; step, stumble, or fall; or remain out of the start position for more than 5 seconds) then this may suggest a concussion.

Any athlete with a suspected concussion should be IMMEDIATELY REMOVED FROM PLAY, urgently assessed medically, should not be left alone and should not drive a motor vehicle.